

On the Road to the Modeling of Resonance UV/vis Spectroscopic Properties

Éric Brémond,^{1,*} Philippe Lainé,¹ François Maurel¹

¹Université de Paris, ITODYS, UMR CNRS 7086, 15 rue J.-A. de Baïf, F-75013 Paris, France

*Corresponding and presenting author: eric.bremond@u-paris.fr

Abstract

Upon a resonant excitation by a specific wavelength, the collectively oscillating electrons in the conduction band of a noble metal nanoparticle, *i.e.* localized surface plasmon resonance (LSPR), prompt a strong light absorption in the UV/vis region. The wavelength coincidence of a chromophore with LSPR leads to an enhanced optical response of the system and could facilitate its study at low concentrations [1].

The state-of-the-art modeling of the resonance UV/vis spectroscopic properties relies on a specific hybrid discrete interaction/quantum model (QM/DIM) which describes the chromophore at TDDFT level, in presence of an electrodynamic treatment of the nanoparticle, including electrostatics and polarization of the atoms [2]. However, prior to applying such a complex model, the nanosystem under investigation has to be carefully chosen. Indeed, a resonant coupling between a large inorganic nanoparticle and a small organic chromophore can only occur in specific conditions, thereby necessitating the design of both nanosized systems. We propose here to investigate the optical properties of a new class of versatile chromophores [3] whose first singlet-singlet absorption energy can be tuned to be in resonance with the surface plasmon of a noble metal nanoparticle. At lower size scale, we further show that the coupling between an organic chromophore and an inorganic metal cluster is a methodological challenge. We thus demonstrate how our last developments in DFT can provide some answers to that specific issue [4].

References

- [1] M. A. El-Sayed *Annu. Rev. Phys. Chem.* 54, 331 (2003).
- [2] J. L. Payton, S. M. Morton, J. E. Moore and L. Jensen *J. Chem. Phys.* 136, 214103 (2012).
- [3] R. Gueret, L. Poulard, M. Oshinowo, J. Chauvin, M. Dahmane, G. Dupeyre, P. P. Lainé, J. Fortage, M.-N. Collomb *ACS Catal.* 8, 3792 (2018).
- [4] É. Brémond, A. J. Pérez-Jiménez, J. C. Sancho-García and C. Adamo *J. Chem. Phys.* 150, 201102 (2019).

Keywords: *resonance UV/vis spectroscopy, carbenium chromophore, surface-enhanced absorption, range-separated exchange density functionals*